

Trial/Project	Sponsor/Funder	Type/Category	Location	Population	Design/Key questions	Status	Status Details
DREAMS	PEPFAR, BMFG, Nike Foundation	Implementation Project	Kenya, South Africa, Swaziland, Uganda, Lesotho, Tanzania, Zambia Zimbabwe	AGYW	Partnership to reduce HIV infections among AGYW; extends beyond health sector to address poverty, gender inequality, sexual violence, lack of education; PrEP implementation component included.	Ongoing	Ongoing.
HPTN 082: Evaluation of daily oral PrEP as a primary prevention strategy for young African women: A Vanguard Study	Wits RHI, HPTN; DAIDS, NIAID, NIMH	Demonstration Project	South Africa; Zimbabwe	Young women (16-25 years)	The goal of this study is to evaluate whether HIV-uninfected sub-Saharan African women ages 16-25 who are at high risk for HIV infection will initiate PrEP and achieve sufficient adherence using scalable adherence support interventions to achieve HIV prevention benefits from this promising biomedical prevention intervention.	Ongoing	Ongoing; started June 2016. Expected completion October 2018.
IMPAACT 2009 (DAIDS ID 30020): Feasibility, Acceptability and Safety of Oral Pre-Exposure Prophylaxis for Primary HIV Prevention during Pregnancy and Breast Feeding in Adolescents and Young Women	International Maternal Pediatric Adolescent AIDS Clinical Trials (IMPAACT) Network	Observational study	Zimbabwe, South Africa, Malawi, Uganda	Adolescent girls and young women aged 16-24 years	Parallel, observational cohort study of HIV-uninfected pregnant adolescents and young women (aged 16-24). The study is designed to characterize adherence over time among women who initiate once daily oral PrEP during pregnancy and continue in the first 6 months following delivery, and to compare pregnancy outcomes among women who take PrEP and women who decline PrEP during the antenatal period.	Planned	Planned; Expected start early-2018.
MTN-034/IPM 045	MTN; IPM; NIH	Phase IIa, Open Label	South Africa; Kenya; Zimbabwe	Adolescent girls and young women aged 16-17 years	Enrolling approximately 300 healthy, HIV-uninfected adolescent females between the ages of 16 - 17 years old (inclusive). Participants will be randomized to product sequence and will use a silicone elastomer vaginal matrix ring (VR) containing 25 mg of dapivirine (DPV) to be replaced each month for a total period of 6 months of use, as well as take oral emtricitabine/tenofovir (FTC/TDF) tablets daily for a 6 month product use period. The primary purpose of this trial is to collect safety and adherence data over the course of study product use. The study will also examine the acceptability of the study products. It is anticipated that this study will take approximately 12 months at each site to enroll the target sample size. Each participant will be followed for approximately 49 weeks.	Planned	Planning underway, expected start early 2018.
PEPFAR	PEPFAR	Implementation Project	South Africa, China, Laos, Thailand, Botswana, Cote d'Ivoire, DRC, Dominican Republic, Ethiopia, Haiti, Kenya, Lesotho, Mozambique, Namibia, Swaziland, Uganda, Ukraine, Vietnam, Tanzania, Zambia, Zimbabwe	At risk individuals	Provides PrEP to adults and adolescents at risk of HIV infection.	Ongoing	Ongoing.
PrEP Operational Research Project	Médecins Sans Frontières	Demonstration Project	Mozambique, Malawi, Zimbabwe	MSM, FSW	Operating in the "corridor," a busy trade route frequented by truckers that runs through Mozambique, Malawi, and Zimbabwe. MSF is providing HIV screening and care to sex workers, drivers, and MSM in the region.	Ongoing	Ongoing; started in 2016.
Sisters Antiretroviral therapy Programme for Prevention of HIV –an Integrated Response (SAPPH-ire)	DFID; UNFPA	Demonstration Project	Zimbabwe	FSW, AGYW	Seeks to enhance HIV treatment and prevention among highway-based sex workers at 7 sites by increasing uptake and frequency of testing, demonstrate acceptability and feasibility of delivering PrEP, maximize retention in care, promote timely initiation of ART for those eligible, and maximize adherence to both ART and PrEP. The evaluation will be in a representative population based sample of 2,800 sex workers, with an estimated 1,000 to 1,500 targeted to take PrEP, after 18 months.	Completed	Completed.