


New CAB PrEP Client or Returning Non-adherent* CAB PrEP Client



Returning Adherent** CAB PrEP Client




TABLE 1: Today's Visit to Initiate/Continue CAB PrEP

Use this table to determine whether to start/continue CAB PrEP, start PEP instead, or defer start of/pause CAB PrEP in a client with a **NEGATIVE 3rd generation rapid diagnostic test (3rd gen RDT) AND clinically assessed as having PEP indication or suspected of acute HIV infection.** All notation of 3rd generation RDT is in accordance with RDTs specified in your national testing algorithm.

New CAB PrEP Clients + Returning CAB PrEP Clients: Non-adherent*

HIGH PEP INDICATION:

- Do **not** initiate or continue CAB PrEP
- Start one-month PEP course
- If available, request expedited RNA PCR on sample collected same day as 3rd gen RDT (prior to PEP) through. **Do not wait for RNA PCR result to start PEP.**
 - If RNA detectable, refer to HIV care and treatment for HIV confirmation
 - If RNA undetectable, complete one-month PEP course
- Post-PEP, re-screen for CAB PrEP, depending on one-month PEP adherence:
 - PEP adherent***: rule-out HIV infection with 3rd gen RDT(s) (Step #1); PEP and AHI assessment (Steps #2 and #3) not needed
 - PEP non-adherent: rule-out HIV infection with Steps 1-3 (3rd gen RDT, PEP assessment, AHI assessment)
 - If CAB PrEP initiated or restarted post-PEP, inject CAB PrEP upon completion of PEP (no break between PEP and PrEP).

HIGH SUSPICION OF AHI:

- Do not initiate or continue CAB PrEP
- Do not start PEP
- If biomedical prevention is desired, PrEP Ring may be offered; do not offer oral PrEP
- If available, request expedited RNA PCR on sample collected same day as 3rd gen RDT (prior to PEP).
 - If RNA detectable, refer to HIV care and treatment for HIV confirmation
 - If RNA undetectable, re-screen for CAB PrEP with Steps 1-3 (3rd gen RDT, PEP assessment, AHI assessment)
- In one-month, re-screen for CAB PrEP with Steps 1-3 (3rd gen RDT, PEP assessment, AHI assessment)

Returning CAB PrEP Clients: Adherent**


- PEP is not indicated among adherent CAB PrEP clients. Continue CAB PrEP regardless of potential exposure or AHI signs and symptoms.

*CAB PrEP Non-adherent: Prior injection was initiation injection 1 and > 2 months ago; prior injection was initiation injection 2 or follow-up and > 3 months ago

**CAB PrEP Adherent: Prior injection was initiation injection 1 and ≤ 2 months ago; or prior injection was initiation injection 2 or follow-up and < 3 months ago

***PEP Adherent: At time of assessment, client reports completion of the one-month PEP regimen

New Oral PrEP Client or Returning Non-adherent* Oral PrEP Client



Returning Adherent** Oral PrEP Client




TABLE 1: Today's Visit to Initiate/Continue Oral PrEP

Use this table to determine whether to start/continue oral PrEP, start PEP instead, or defer start of/pause oral PrEP in a client with a **NEGATIVE 3rd generation rapid diagnostic test (3rd gen RDT) AND clinically assessed as having PEP indication or suspected of acute HIV infection**. All notation of 3rd generation RDT is in accordance with RDTs specified in your national testing algorithm.


New Oral PrEP Clients + Returning Oral PrEP Clients: Non-adherent*	
HIGH PEP INDICATION:	
<ul style="list-style-type: none"> Do <u>not</u> initiate or continue oral PrEP Start one-month PEP course Post-PEP, re-screen for oral PrEP, depending on one-month PEP adherence <ul style="list-style-type: none"> PEP adherent***: rule-out HIV infection with 3rd gen RDT (Step #1); PEP and AHI assessment (Steps #2 and #3) not needed PEP non-adherent: rule-out HIV infection with Steps 1-3 (3rd gen RDT, PEP assessment, AHI assessment) If oral PrEP initiated or restarted post-PEP, dispense oral PrEP upon completion of PEP (no break between PEP and PrEP) 	
HIGH SUSPICION OF AHI:	
<ul style="list-style-type: none"> Do <u>not</u> initiate or continue oral PrEP Do <u>not</u> start PEP If biomedical prevention desired, PrEP ring may be offered; do <u>not</u> offer oral PrEP In one-month, re-screen for oral PrEP with Steps 1-3 (3rd gen RDT, PEP assessment, AHI assessment) 	
Returning Oral PrEP Clients: Adherent**	
<ul style="list-style-type: none"> PEP is not indicated among adherent oral PrEP clients. Continue oral PrEP regardless of potential exposure or AHI signs and symptoms. 	

*Oral PrEP Non-adherent: 4+ doses missed each week in week before and week after each day of sex

**Oral PrEP Adherent: ≤ 3 doses missed each week in week before and week after each day of sex

***PEP Adherent: At time of assessment, client reports completion of the one-month PEP regimen

New PrEP Ring Client or Returning Non-adherent* PrEP Ring Client



Returning Adherent** PrEP Ring Client




TABLE 1: Today's Visit to Initiate/Continue PrEP Ring

Use this table to determine whether to start/continue PrEP ring or start PEP instead in a client with a **NEGATIVE 3rd generation rapid diagnostic test (3rd gen RDT) AND clinically assessed as having PEP indication or suspected of acute HIV infection.** All notation of 3rd generation RDT is in accordance with RDTs specified in your national testing algorithm.

New PrEP Ring Clients + Returning PrEP Ring Clients: Non-adherent*

HIGH PEP INDICATION:

- Do not initiate or continue PrEP ring
- Start one-month PEP course
- Post-PEP, re-screen for PrEP ring, depending on one-month PEP adherence
 - ▶ PEP adherent***: rule-out HIV infection w/ 3rd gen RDT(s) (Step #1); PEP and AHI assessment (Steps #2 and #3) not needed
 - ▶ PEP non-adherent: rule-out HIV infection with Steps 1-3 (3rd gen RDT, PEP assessment, AHI assessment)
 - ▶ If PrEP ring initiated or restarted post-PEP, dispense PrEP ring (no break between PEP and PrEP)

HIGH SUSPICION OF AHI:

- PrEP ring use is permissible despite AHI suspicions: start or continue PrEP ring despite AHI suspicions
- Do not start PEP
 - ▶ Continue with HIV re-testing by 3rd Gen RDT as per scheduled visits (at least quarterly)

Returning PrEP Ring Clients: Adherent**

- Provide option of HIV prevention approach based on likelihood and recency of potential HIV exposure:
 - ▶ If particularly high likelihood of exposure within 72 hours, offer the option to stop PrEP ring and switch to one-month course of PEP
 - ▶ Continue PrEP ring if lesser likelihood or beyond 72 hours, or per client preference

*PrEP Ring Non-adherent: Ring not continuously in place the day before and day of vaginal sex; any reported anal sex without a condom should be considered PrEP ring non-adherent

**PrEP Ring Adherent: Ring in place continuously day before and day of sex for each day of vaginal sex

***PEP Adherent: At time of assessment, client reports completion of the one-month PEP regimen