Course: Providing the PrEP ring to Pregnant and Breastfeeding People — Training Pre- and Post- Test Answer Key

Instructions: This assessment includes 10 questions and should take around 10 minutes. Choose one answer per question, unless the question specifies to 'select all that apply.'

- 1. Why is it important to offer PrEP method choice to pregnant and breastfeeding people?
 - a. People are at increased risk of HIV acquisition during pregnancy and breastfeeding.
 - b. PrEP products are more effective in pregnant clients than in clients who are not pregnant.
 - c. People are more likely to agree to take PrEP during pregnancy than they are at other times.
- 2. Which common medications prescribed during pregnancy can have unwanted interactions with the PrEP ring?
 - a. PrEP ring may interact with antenatal medications such as iron and folic acid tablets.
 - b. PrEP ring may interact with malaria treatments, including sulfadoxine-pyrimethamine.
 - c. The medications used in the PrEP ring have no known drug interactions with the most commonly prescribed pregnancy medications.
- 3. Which of the following clients are good candidates for the PrEP ring? Select all that apply.
 - a. An ANC client who is using oral PrEP, but is having difficulty remembering to take pills every day.
 - b. A client who is already using the ring but becomes pregnant.
 - c. A client with exposure to HIV in the past 72 hours.
 - d. A client who is one day postpartum and actively bleeding.
- 4. Which of the following are correct statements when counseling pregnant and breastfeeding people on the use of the PrEP ring? **Select all that apply.**
 - a. PrEP ring use during pregnancy has not been shown to cause babies to be too big or too small.
 - b. PrEP ring use may impact a person's ability to get pregnant in the future.
 - c. PrEP ring has not been shown to affect a mother's milk production or the taste or quality of breast milk.
- 5. Which of the following are contraindications to starting PrEP ring? Select all that apply.
 - a. Breastfeeding
 - b. Suspected or confirmed intrauterine infection following a miscarriage
 - c. Active labor at any gestational age







- 6. What are some ways PrEP providers could address pregnant client concerns regarding potential side effects they may be experiencing using PrEP ring? Select all that apply.
 - a. Counseling on common side effects experienced during pregnancy to rule out pregnancy-related side effects
 - b. History taking and targeted physical examination
 - c. Client-centered plan, including planning for follow-up and future evaluation to see if concerns have resolved
 - d. All of the above
- 7. Which of the following is **not** true regarding removing and inserting the PrEP ring?
 - a. The PrEP ring should be removed before sex as it will likely be felt by sexual partner and interfere with sex.
 - b. The PrEP ring should be removed before delivering the baby, but can be re-inserted after delivering the baby, ideally after the uterus has returned to its pre-pregnancy size and bleeding has diminished.
 - c. Clients can insert, remove, and replace the PrEP ring themselves each month.
- 8. True or False: When a pregnant or breastfeeding person who is using the PrEP ring reports a possible side effect, the PrEP ring should immediately be paused while the cause of the symptom is evaluated.
 - a. True
 - b. False
- 9. True or False: HIV testing services are not required while a pregnant or breastfeeding person is using the PrEP ring.
 - a. True
 - b. False
- 10. What does the acronym LIVES stand for when defining first-line support goals for survivors of violence?
 - a. Listen with empathy, Inquire about needs and concerns, Validate the survivor's experience, Enhance safety, and Support the survivor
 - b. Listen with empathy, Involve the survivor in next steps, View previous records and history, Empathize with the survivor, and Support the survivor
 - c. Look up survivor records and history, Include the survivor and partner in decisions, Validate the survivor's story, Education the survivor about options, and Support the survivor