





JOB AID: ASSESSMENT FOR PREP CLINICAL ELIGIBILITY AND CONTRAINDICATIONS

Assess the following to determine if a client is clinically eligible for PrEP:	Clinical Contraindication or Special Consideration			
	TDF-Based Oral PrEP 	DVR 	CAB-LA 	LEN 
1) HIV Status	If HIV test results are positive or inconclusive, clients should be referred for further testing/treatment.	If HIV test results are positive or inconclusive, clients should be referred for further testing/treatment.	If HIV test results are positive or inconclusive, clients should be referred for further testing/treatment.	If HIV test results are positive or inconclusive, clients should be referred for further testing/treatment.
2) Post-Exposure Prophylaxis (PEP) Indication	If PEP is highly indicated, start client on 1 month course of PEP and defer/pause PrEP.	If PEP is highly indicated, start client on 1 month course of PEP and defer/pause PrEP.	If PEP is highly indicated, start client on 1 month course of PEP and defer/pause PrEP.	If PEP is highly indicated, start client on 1 month course of PEP and defer/pause PrEP.
3) Suspicion of Acute HIV Infection (AHI)	If high suspicion of AHI, defer/pause PrEP. Advise client to return in 1 month for re-testing	If high suspicion of AHI, proceed with eligibility screening and re-test in 1 month. DVR may be used despite suspicion of AHI	If high suspicion of AHI, defer/pause PrEP. Advise client to return in 1 month for re-testing	If high suspicion of AHI, defer/pause PrEP. Advise client to return in 1 month for re-testing
4) Possible Medication Interactions	<p><u>Contraindications:</u></p> <p>TDF-based oral PrEP is not contraindicated with most commonly used medicines</p> <p><u>Considerations:</u></p> <p>If client is taking estradiol-based hormones: only use TDF-based dosing regimen/schedule recommended for cis-gender women, transgender women taking gender-affirming hormones, and transgender men</p>	<p><u>Contraindications:</u></p> <p>Do not use in combination with contraceptive vaginal rings or diaphragms; avoid use with vaginally administered metronidazole or clindamycin</p> <p><u>Considerations:</u></p> <p>Concomitant use of vaginally administered clotrimazole should be with caution</p> <p>Offer condoms if concomitantly used with vaginally administered miconazole</p>	<p><u>Contraindications:</u></p> <p>Do not use in combination with:</p> <ul style="list-style-type: none">• rifampin• oxcarbazepine• rifapentine• phenobarbital• carbamazepine• phenytoin <p><u>Considerations:</u></p> <p>Use in combination with rifabutin may require a dose adjustment, and prescribers should review the regulatory labels of both medications when being prescribed for concomitant use</p>	<p><u>Contraindications:</u></p> <p>There are no known medications contraindicated with LEN</p> <p><u>Considerations:</u></p> <p>Use of any of the following medications concomitantly with LEN or in the months after LEN discontinuation may require a dose adjustment (to the dose of either LEN or the other medication):</p> <ul style="list-style-type: none">• Rifabutin• Rifampicin• Rifapentine• Carbamazepine• Phenobarbital• Phenytoin• Ketamine• Avanafil• Sildenafil• Tadalafil• Vardenafil
5) Possible Medication Allergies	If client has had previous allergic reaction to TDF, FTC or 3TC-based medications, recommend an alternative PrEP product	If client has had previous allergic reaction to DVR, recommend an alternative PrEP product	If client has had previous allergic reaction to cabotegravir, recommend an alternative PrEP product	If client has had previous allergic reaction to lenacapavir, recommend an alternative PrEP product
6) Relevant comorbidities	If client has impaired Renal Function (CrCl< 60 ml/min or eGFR 60 ml/min per 1.73 m ²), recommend an alternative PrEP product	If client has severe vaginal health problems, treat before starting or continuing with DVR.	If client is known to have impaired liver function (i.e. chronic liver disease or acute hepatitis) recommend an alternative PrEP product	There are no known comorbidities that require assessment
	<i>Note: renal function testing optional if client <49 years and free of renal comorbidities</i>	<i>Note: speculum examination is not required</i>	<i>Note: testing for viral hepatitis or liver function is not required prior to starting CAB-LA</i>	