


# JOB AID: TRANSITIONING BETWEEN PREP PRODUCTS AND OVERLAPPING USE CONSIDERATIONS TO ENSURE PROTECTION

## TRANSITIONING FROM ORAL TDF-BASED PREP


TO DVR

Clients should continue using TDF-based oral PrEP for **7 days** after insertion of DVR




TO CAB-LA

Clients should continue using TDF-based oral PrEP for **7 days** after CAB-LA INITIATION INJECTION #1



TO LEN


Clients should continue using TDF-based oral PrEP for **2 days** after LEN INITIATION REGIMEN



## TRANSITIONING FROM DAPIVIRINE VAGINAL RING (DVR)


TO ORAL PREP

Clients should continue using DVR for **7 days** after starting TDF-based oral PrEP




TO CAB-LA

Clients should continue using DVR for **7 days** after CAB-LA INITIATION INJECTION #1



TO LEN

Clients should continue wearing DVR throughout **both days** of receiving the INITIATION REGIMEN to start LEN (all doses)



# TRANSITIONING FROM CAB-LA

TO ORAL PREP

If last injection was INITIATION INJECTION #1, client should begin oral TDF-based PrEP **1 month** after last injection

CAB-LA INITIATION INJECTION #1



START ORAL PREP




Future oral PrEP use

If last injection was INITIATION INJECTION #2 or FOLLOW-UP INJECTION, client should begin oral TDF-based PrEP **2 months** after last injection

Prior CAB-LA use

CAB-LA INITIATION INJECTION #2 or FOLLOW UP INJECTION



START ORAL PREP



Future oral PrEP use

TO DVR

If last injection was INITIATION INJECTION #1, client should insert DVR **1 month** after last injection

CAB-LA INITIATION INJECTION #1



INSERT DVR



Future DVR use

If last injection was INITIATION INJECTION #2 or FOLLOW-UP INJECTION, client should insert DVR **2 months** after last injection

Prior CAB-LA use

CAB-LA INITIATION INJECTION #2 or FOLLOW UP INJECTION



INSERT DVR



Future DVR use

TO LEN

If last CAB-LA injection was INITIATION INJECTION #1, client should start LEN INITIATION REGIMEN **1 month** after last CAB-LA injection

CAB-LA INITIATION INJECTION #1



LEN INITIATION REGIMEN



Future LEN Injections

If last CAB-LA injection was INITIATION INJECTION #2 or FOLLOW-UP INJECTION, client should start LEN INITIATION REGIMEN **2 months** after last CAB-LA injection

Prior CAB-LA use

CAB-LA INITIATION INJECTION #2 or FOLLOW UP INJECTION



LEN INITIATION REGIMEN



Future LEN Injections


# TRANSITIONING FROM LEN

TO ORAL PREP

Clients should start start TDF-based oral PrEP **6 months** after the last LEN injections were administered, regardless of the type of LEN injections last administered

Prior LEN use

LEN INJECTIONS



START ORAL PREP




Future oral PrEP use

TO DVR

Clients should insert DVR **6 months** after the last LEN injections were administered, regardless of the type of LEN injections last administered

Prior LEN use

LEN INJECTIONS



INSERT DVR




Future DVR use

TO CAB-LA

Clients should start CAB-LA **6 months** after the last LEN injections were administered, regardless of the type of LEN injections last administered

Prior LEN use

LEN INJECTIONS



CAB-LA INIT INJ#1



Future CAB-LA Injections