


CAB-LA Missed Visits Scheduling Wheel

This wheel is used to support providers in clinically managing clients who miss a long-acting cabotegravir (CAB-LA) injection. To print and assemble this wheel, please follow these instructions:

- Print in color on heavy card stock / artboard, or plastic
 - (Pages 2 & 3 – Outer wheel front & back double sided)
- Diecut to shape
- Matte laminate
- Assemble with a rivet through the center to join rotating Outer and Inner wheels
 - Suggested minimum Outer wheel diameter of 5 inches / 127mm, depending on local preference





CAB-LA MISSED INJECTION WHEEL

Period A:

Resume with either INITIATION INJECTION #2 or FOLLOW-UP INJECTION today if clinically eligible; schedule next FOLLOW-UP INJECTION in 2 months

Period B:

Restart CAB-LA with INITIATION INJECTION #1 today if clinically eligible; schedule INITIATION INJECTION #2 in 1 month



Before resuming or restarting injections, assess for PEP and AHI as part of determining clinical eligibility to continue CAB-LA use, as follows:

Ensure PEP is not indicated. PEP indication based on possible HIV exposure past 72 hours **AND** significance of delay in returning for CAB-LA reinjection (use deviation). If high indication for PEP, consider starting 1-month course of PEP today instead of resuming CAB-LA use.

Ensure AHI is not suspected. AHI suspicion based on signs and symptoms (past 2 weeks and possible HIV exposures in the past 1 month) **AND** significance of delay in returning for CAB-LA reinjection. If high suspicion of AHI, consider pausing CAB-LA use pending HIV retesting in 1 month; advise alternative HIV prevention strategies pending retesting. Note: signs/symptoms may be absent despite AHI due to possibilities of LEVI syndrome.

